

International Encyclopedia of Rehabilitation

Copyright © 2010 by the Center for International Rehabilitation Research Information and Exchange (CIRRIE).

All rights reserved. No part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system without the prior written permission of the publisher, except as permitted under the United States Copyright Act of 1976.

Center for International Rehabilitation Research Information and Exchange (CIRRIE)
515 Kimball Tower
University at Buffalo, The State University of New York
Buffalo, NY 14214
E-mail: ub-cirrie@buffalo.edu
Web: <http://cirrie.buffalo.edu>

This publication of the Center for International Rehabilitation Research Information and Exchange is supported by funds received from the National Institute on Disability and Rehabilitation Research of the U.S. Department of Education under grant number H133A050008. The opinions contained in this publication are those of the authors and do not necessarily reflect those of CIRRIE or the Department of Education.

Prehension

Luciane Sande

Department of Physiotherapy, Federal University of Sao Carlos

Summary

The important function of prehension is explored in different points, including its development, which helps the baby in the learning process. Because of the multiple degrees of freedom, the hand can be molded in many meaningful ways. These patterns are presented according to the literature evolution. Motor prehension control shows how the central nervous leads with the several prehension conditions. Also, its evaluation can help in many diagnosis processes. Related disorders with some physical interventions are presented in the next section.

Definition and function of prehension

The human hand is a complex instrument, that has multiple aims. In a physiologic view, the hand represents the effector extremity of the arm, as it can be in the most favorable position to execute specific functions (Lehmkuhl; Smith 1989). The functional capacities of the hand as a perfect locomotor system organ have great influence in the social and creative efficiency of humans. Only the hand is capable of making minucious distinctions about external characteristics because it combines strength and accuracy (Catovic et al 1991).

Using the independent movements of the several segments of fingers and hand we can hold, maintain, release, take the hand to objects or handle them and move them in all directions. In all these actions there is: 1) the independence of thumb movements and indicator among themselves and in relation to the other fingers, 2) the fact that all fingers movements and wrist run independently of the arm segments positions, 3) independent movements of the interphalangeal and metacarpophalangeal joints (Brandão 1984).

Although the hand has multiple functions, the essential function is prehension. Such capacity appears from the tweezers lobster to the hand of the monkey, but in humans prehension reached its highest degree of functionality. This is due to a particular provision of thumb, which may object to all other fingers (Kapandji 1980).

Human manipulative tasks are carried out by hand movements, that frequently involve forceful movements and prehension. According to Putz-Andersson (1988) the movement of the thumb causes the hand to be aligned with the forearm, resulting in a slight ulnar deviation of the wrist.

Prehension development

The maturity of the different parts of the central nervous and musculoskeletal systems are associated with the mature patterns of prehension acquired in the first years of the life (Cook; Woollacott 1995). As sensory-motor coordination develops, the adult characteristics of prehension are also imprinted in the motor memory of the child. (Brandão 1984). Hand-eye coordination helps in eye fixation, which means a good eye stabilization, and it is also presented during the movement of a moving target. Like the motor components, the sensorial

contributions to prehension are essentials to a good and accurate performance (Cook; Woollacott 1995).

Babies use their hand to learn and explore the environment and indirectly they develop the concepts of spatial dimensions of objects and other physical properties, like shape and texture (Imrhan 1989). After infancy, the hand assumes more specific roles, and can be used to access and shape the environment.

Prehension patterns and behaviour

Prehension patterns have not been clearly defined. According to Bendz (1974) there is much confusion generated by inadequate synonymy. The following are some classifications of prehension patterns in the chronological order in which they were presented in the literature.

The beginning

Schlesinger in 1919 (apud Jain, Hennedy and Carus 1985) related six basic patterns of prehension: cylindrical, in point, on hook, palm, spherical and lateral. Later in 1947, of the University of Berkeley, California (Jain, Hennedy and Carus 1985), a study using kinematography identified three basic patterns of the six previous: palm, in point and lateral. Such studies were added to others (Griffiths 1943; Napier 1956; Taylor; Schwarzs 1955 (apud Lehmkuhl; Smith 1989) Rash; Burke 1977 (apud Maluf 1986) and Lehmkuhl; Smith (1989) which used the shape of the object as the parameter to the classification.

Criteria for functional classification

Classification based on action

Alongside the classifications described, Napier 1956 (apud Maluf 1986) found that the movements of prehension should not be determined, only by the shape of the object, but also by the purpose of the action. Based on this finding a dynamic and functional aspect of prehension was described, with two basic factors: strength and precision. In the strength prehension, the object is pressed against the palm of the hand for the generation of force by the fingers and thumb. In the precision prehension, the object is manipulated between the thumb and the fingertips in a fine movement without the involvement of the palm.

In 1974, Bendz completed the system of Napier (apud Maluf 1986), proposed in 1956, including the flexor and extensor prehension.

Classification based on the contact area: hand-object

In 1980 Kapandji, taking into account the areas of contact between the object and the fingers and/or the hand, described the following types of grip: opposition terminal prehension, opposition subterminal prehension, opposition subterminal-lateral prehension, palm prehension in “full hand”, palm-digit prehension, interdigit lateral prehension.

Again Kapandji in 1987 (apud Pachioni, Loureiro and Fonseca 1992), improved the classification and grouped prehension in this way: 1) digitals, bidigitals and pluridigitals; 2)

palm; 3) centrated. In another group, the prehensions with weight and actions prehensions were included.

Although there have been changes over time, there is still much confusion in the ranks. There is also some overlap between the patterns of the different classifications. Moreover, there seems to be a trend in the current literature to standardize and group the prehensions again in three forms already recommended by the study of Berkeley University in 1947 (apud Jain, Henedy and Carus, 1985) and by Taylor; Schwatz 1955 (apud Lehmkuhl; Smith 1989). That is, although the systems have become increasingly complex and detailed, the forms most studied, probably because they are the most used, are the three basic forms of grip, namely: lateral, palm and pulp or edge (mainly between thumb and second finger)(Ager, Olivett and Johnson 1984; Mahiowetz 1985; Imrhan 1989; Imrhan 1992; Fernandez et al 1992; Imrhan; Rahman 1995).

Motor control of prehension

Cole and Abbs (1988) and Flanagan and Wing (1995) identified a parallel modulation of prehensile strength and strength to lift objects, independant of surface texture and frequency of movement. In the same way, Westling and Johansson (1988) described motor commands which varied with the load being handled. In this way it is believed that daily practice can lead to an adaptation of these commands, a phenomenon described as adaptive memory. Thus each time the weight is altered, the command is corrected (Sangole; Levin 2007).

It is believed that the perception of muscular contraction can be codified in the memory by two mechanisms: one from the pressure of the reaction occurring at the surface of the palm and fingers, which stimulates the mechanoreceptors of the region, and the other a kinesthetic model, generated from the muscular tension of the phalanx flexor muscles (Lowe, 1995). Thus, the sensory inputs reveal important functions in the control of movements, such as the modulation of the output of the muscles, optimizing the strength to be used (Cook; Woollacott, 1995).

Various individual factors have been recognized as capable of affecting prehensile strength in normal individuals, including age (Mathiowetz et al 1985; Imrhan 1989), sex and body posture (Catovic et al 1991; Fernandez et al 1992; Lamoreaux; Hoffer 1995). Other factors such as hand size and arm length, have also been associated with prehensile strength, although they are more controversial.

Also the posture used during prehension must be considered. The orthostatic position compared with seat and supine postures seems to facilitate higher strength levels be achieved in the prehension movement (Su et al 1994; Catovic et al 1989; 1991).

Evaluation of hand grip strength

Pereira et al (2001) said that grip strength can be used as an indicator of coordination development and helping in the diagnosis of neurological disfunctions related to the motor learning and to perception and also in the identification of arm, forearm and hand

pathology, in the control of the rehabilitation process and in the determination of supportable patterns in the application or sustentation of loads.

Evaluating prehension strength in children of 7 to 14 years, Esteves et al (2005) reported that several factors can influence strength production: age, laterality, practice in different physical activities, size of the grip, sex, height and body mass. These authors also emphasized that the values of prehension strength are maximal in the adult phase, about 25-35 years. After this period, there is a gradual decline.

In the same way, Barbosa et al (2006) exploring the relation between strength prehension and the nutritional state in elderly persons, concluded that there is a positive correlation between them. There are modifications in the nutritional state and muscle strength with the advance of age; the effects of the nutritional state on the muscle strength are specific to the nutritional indicator, to the sex and to the age group.

Some authors identified reduction in maximum strength and prehensile capacity in the presence of pain and musculoskeletal disorders (Smith; Bengt 1985; Janda et al 1987; Niebuhr; Marion 1987).

Musculoskeletal disorders related to prehension

According to Putz-Andersson (1988) non-neutral postures of the wrist combined with forceful and repetitive movements are important risk factors for triggering upper limb work-related musculoskeletal disorders (WRMDs).

Muscle, tendon and nerve respond to dynamically varying local forces. The stresses generated under these situations were reviewed by Hagberg et al (1995). According to this review, both external (force exertion) and internal forces (from stretching or compression of the median nerve at the wrist) are hypothesized to be part of the mechanisms leading to WRMDs. Chronic disorders usually present more structural tissue damage. A decrease in the capacity to exert strength is one of the outcomes of tissue damage. Therefore, measurements of the hand grip strength are used as an indicator of impairment of the hand function.

Basic rehabilitation interventions

In abnormal neuromotor development we can find children without the adequate and necessary sensorimotor experiences due to abnormal primitive reflexes and abnormal muscle tone. Consequently, in a special treatment, using recreational activities, including the handling of various types of toys (textures, shapes, weights, sizes), we must inhibit these reflexes to facilitate normal muscle tone. Moreover, to prepare the ground for the coordination of grip and precise handling, techniques of facilitation should be employed within the sequential development process (Duff; Charles 2004).

Similarly, when an adult suffers brain damage, the grip can also be damaged. The use of compensatory strategies for prehension may be related to the degree of motor impairment: severely to moderately impaired subjects recruited new degrees of freedom to compensate for motor deficits, while mildly impaired subjects tended to employ healthy movement

patterns (Cirstea; Levin 2000). Through physiotherapy these patients may have an improvement in its grip with exercises that include strength, elongation, facilitation of weak and little active muscles and also the use of mental practice. In addition, placements and adequate incentives for the intense use of common prehensions in activities of daily living can be introduced. There is also the so-called Constraint induced therapy, which is based on restriction of the less affected arm allowing concentrated facilitation of the more-affected arm (Wolf et al 2008).

Diseases or injuries affecting the spinal cord, peripheral nerves or even the osteomyoarticular system can also compromise the use of prehensions. Again, several classes of exercises and even the use of functional electrical stimulation can result in improvement in the strength and accuracy of the various forms of grip. Some novel interventions can also be suggested: massed practice (repetitive activity-based training) and somatosensory stimulation (prolonged peripheral nerve electrical stimulation at submotor threshold intensity)(Beekhuizen 2005).

As commented before, a great number of lesions are related to the work, so many modifications have to be considered: repetition of the same task must be avoided, a rest period must be respected, the exercises practiced, specially elongations must be intense. In some cases the subjects have to change their work activity and also be submitted to a surgery with a long rehabilitation period. (Spielholz et al 2001).

References

- Ager CL, Olivett BL, Johnson CL. 1984. Grasp and Pinch Strength in Children 5 to 12 Years old. *American Journal of Occupational Therapy* 38(2):107-13.
- Barbosa AR, Souza JMP, Lebrão ML, Marucci MFN. 2006. Relação entre estado nutricional e força de preensão manual em idosos do município de São Paulo, Brasil: dados da pesquisa SABE. *Revista Brasileira de Cineantropometria and Desempenho Humano* 8(1):37-44.
- Beekhuizen KS. 2005. New perspectives on improving upper extremity function after spinal cord injury. *Journal of Neurologic Physical Therapy* 29(3):157-62.
- Bendz P. 1974. Systematization of the grip of the hand in relation to finger motor systems. A kinesiologic study using a new method for recording finger joint motions. *Scandinavian Journal of Rehabilitation Medicine* 6(4):58-65.
- Brandão JS. 1984. *Desenvolvimento Psicomotor da Mão*. RJ: Enelivros.
- Catovic A, et al. 1989. A comparative investigation of the influence of certain arm positions on hand pinch grips in the standing and sitting positions of dentists. *Applied Ergonomics* 20(2):109-114.

- Catovic E, Catovic A, Kraljevic K, Muftic O. 1991. The influence of arm position on the pinch grip strength of female dentists in standing and sitting positions. *Applied Ergonomics* 22(3):163-166.
- Cirstea MC, Levin MF. 2000. Compensatory strategies for reaching in stroke. *Brain*.123 (Pt 5):940-53.
- Cole KJ, Abbs JH. 1988. Grip force adjustments evoked by load force perturbations of a grasped object. *Journal of Neurophysiology* 60(4):1513-1522.
- Cook AS, Woollacott MH. 1995. *Motor control: Theory and Practical Applications*. Williams and Wilkins.
- Duff SV, Charles J. 2004. Enhancing prehension in infants and children: fostering neuromotor strategies. *Physical and Occupational Therapy in Pediatrics* 24(1-2):129-72.
- Esteves AC, Reis DC, Caldeira RM, Leite RM, Moro ARP, Borges NG. 2005. Força de preensão, lateralidade, sexo e características antropométricas da mão de crianças em idade escolar. *Revista Brasileira de Cineantropometria and Desempenho Humano* 7(2):69-75.
- Fernandez JE, Dahalan JB, Halern CA, Fredericks TK. 1992. The effect of deviated wrist posture on pinch strength for females. In: *Advances in Industrial Ergonomics and Safety, IV*. Taylor & Francis. p 693-700.
- Flanagan JR, Wing AM. 1995. The stability of precision grip forces during cyclic arm movements with a hand-held load. *Experimental Brain Research* 105:455-464.
- Hagberg M, Silverstein B, Wells R, et al. 1995. *Work related musculoskeletal disorders (WMSDs). A reference book for prevention*. Ed.by I. Kuorinka and L. Forcier. Taylor & Francis.
- Imrhan SN. 1989. Trends in finger pinch strength in children, adults, and elderly. *Human Factors* 31(6):689-701.
- Imrhan SN, Sundararajan K. 1992. An investigation of finger pull strengths. *Ergonomics* 35(3):289-299.
- Imrhan SN, Rahman R. 1995. The effects of pinch width on pinch strengths of adult males using realistic pinch-handle coupling. *International Journal of Industrial Ergonomics* 16:123-134.
- Jain AS, Henedy JA, Carus DA. 1985. Clinical assessment of hand strength using a microcomputer. *Journal of Hand Surgery* 10-B:3, 315-318.
- Janda, DH, et al. 1987. Objective evaluation of grip strength. *Journal of Occupational Medicine* 29(7):569-571.

- Kapandji IA. 1980. Fisiologia Articular: esquemas comentados de mecânica humana, 4th ed. São Paulo: Manole.
- Lamoreaux L, Hoffer MM. 1995. The effect of wrist deviation on grip and pinch strength. *Clinical Orthopedics and Related Research* 314:152-155.
- Lehmkuhl L, Smith L. 1989. Clinical Kinesiology from Brunnstrom. 4th ed. São Paulo: Manole. (in Portuguese).
- Lowe B. 1995. Repeatability in magnitude production in isometric, hand grip estimation: a working -memory approach. *Perceptual and Motor Skills* 180:659-667.
- Maluf AS. 1986. Estudo eletromiográfico dos músculos abdutor curto e oponente do polegar em movimentos livres, contra-resistidos e de garra. Piracicaba: Faculdade de Odontologia da UNICAMP. (dissertação).
- Mathiowetz V, Kashman N, Volland G, et al. 1985. Grip and Pinch Strength: Normative Data for Adults. *Archives of Physical Medicine in Rehabilitation* 66: 69-74.
- Niebuhr B, Marion R. 1987. Detecting Sincerity of effort when measuring grip strength. *Americal Journal of Physical Medicine* 66(1):16-23.
- Pachioni CA, Loureiro D, Fonseca MC. 1992. Anatomia funcional e reabilitação nas lesões dos tendões flexores e extensores da mão. Ribeirão Preto: Faculdade de Medicina da USP. (trabalho de especialização).
- Pereira HS, Landgren M, Gillberg C, Forsberg H. 2001. Parametric control of fingertip forces during precision grip lifts un children with DCD (developmental coordination disorder) and DAMP (deficits in attention motor control e perception). *Neuropsychologia* 39:478-488.
- Putz-Andersson V. 1988. Cumulative trauma disorders: a manual for musculoskeletal disease of upper limbs. Taylor & Francis: London.
- Sangole AP, Levin MF. 2007. A new perspective in the understanding of hand dysfunction following neurological injury. *Topics in Stroke Rehabilitation* 14(3):80-94.
- Smith RO, Bengt MW. 1985. Pinch and grasp strength: standardization of terminology and protocol. *American Journal of Occupational Therapy* 39(8):531-535.
- Spielholz P, Silverstein B, Morgan M, Checkoway H, Kaufman J. 2001. Comparison of self-report, video observation and direct measurement methods for upper extremity musculoskeletal disorder physical risk factors. *Ergonomics* 15;44(6):588-613.
- Su CY, et al. 1994. Grip Strenght in different Positions of elbow and shoulder. *Archives of Physical Medicine in Rehabilitation* 75:812-5.

- Westling G, Johansson RS. 1998. Coordinated isometric muscle commands adequately and erroneously programmed for the weight during lifting task with precision grip. *Experimental Brain Research* 71:59-71.
- Wolf SL, Winstein CJ, Miller JP, Thompson PA, Taub E, Uswatte G, Morris D, Blanton S, Nichols-Larsen D, Clark PC. 2008. Retention of upper limb function in stroke survivors who have received constraint-induced movement therapy: the EXCITE randomised trial. *Lancet Neurology* 7(1):33-40.