

International Encyclopedia of Rehabilitation

Copyright © 2010 by the Center for International Rehabilitation Research Information and Exchange (CIRRIE).

All rights reserved. No part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system without the prior written permission of the publisher, except as permitted under the United States Copyright Act of 1976.

Center for International Rehabilitation Research Information and Exchange (CIRRIE)
515 Kimball Tower
University at Buffalo, The State University of New York
Buffalo, NY 14214
E-mail: ub-cirrie@buffalo.edu
Web: <http://cirrie.buffalo.edu>

This publication of the Center for International Rehabilitation Research Information and Exchange is supported by funds received from the National Institute on Disability and Rehabilitation Research of the U.S. Department of Education under grant number H133A050008. The opinions contained in this publication are those of the authors and do not necessarily reflect those of CIRRIE or the Department of Education.

Bone Loss in Spinal Cord Injury and Multiple Sclerosis

Yannis Dionyssiotis

Physical and Social Rehabilitation Center, Amyntæo, Florina, Greece

yannis_dionyssiotis@hotmail.com

Financial

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

Introduction

Osteoporosis is characterized by low bone mass and destruction of the microarchitecture of bone tissue, resulting in increased bone fragility and susceptibility to fractures (NIH 2001). The clinical usefulness of T-score at disabled people on the recognition of people with low BMD remains unclear according to ranking system of the World Health Organization (WHO 1994). Despite the increased number of risk factors in people with disabilities no guidelines are available on BMD measurements; so it would be more appropriate to use the term low bone mass instead of osteoporosis or osteopenia and also take into account the Z-score obtained from the measurement of bone densitometry which is the number of standard deviations above or below that normally expected for someone of similar age, sex, weight and race in question. In disabilities there are differences among subjects according to the type of lesion; complete (an absence of sensory or motor function below the neurological level, including the lowest sacral segment) vs. incomplete lesions (partial preservation of motor and/or sensory function below the neurological level, including the lowest sacral segment), the neurological level of lesion, the progress of the disease (i.e in complete spinal cord injury vs. progressive multiple sclerosis), the residual mobility, walking-standing ability, functionality and degree of spasticity.

Physiopathology of bone loss in spinal cord injury

Bone loss in spinal cord injury (SCI) is a multifactorial disease in acute and chronic phase and can be enhanced by the lack of weight bearing, muscular tension on bone or other neural factors associated with the injury. Moreover, differentiation of the sympathetic nervous system after SCI is leading to venous and capillary vascular stasis. Some additional non-mechanical factors to stimulate bone loss include poor nutritional adequacy, gonadal changes and other endocrine disorders (Chantraine 1978; Chantraine and others 1979b; Jiang and others 2007; Maimoun and others 2006).

A reduction in bone density about 4% per month in the first year was estimated in areas rich in trabecular bone, such as the tibial and femoral epiphyses and about 2% per month in areas with predominantly compact bone (Wilmet and others 1995). In studies with peripheral quantitative computed tomography (p QCT) in paraplegics bone loss in the epiphyses was 50% in the femur and 60% in the tibia, while in the

diaphyses of these bones was 35% and 25%, respectively, meaning that bone loss in the epiphyses almost doubled the loss in the diaphyses (Figure 1). According to bone loss there are some interesting features; demineralization in spinal cord injured subjects is area dependent, occurs exclusively in the areas below the level of injury (Dauty and others 2000), affecting mainly paralyzed extremities and increasing from proximal to distal regions i.e in paraplegics weight bearing skeleton regions, as the distal end of femur and proximal tibia, which are rich in cancellous bone, while region of the diaphysis of the femur and tibia, rich in cortical bone is reserved (Eser and others 2004; Kiratli and others 2000). Moreover, bone loss between trabecular and cortical bone compartment differs in mechanism, i.e in the epiphyses is due to decrease in trabecular but in diaphysis cortical bone is maintained and bone is lost through endocortical resorption by reducing cortical wall thickness (Dionyssiatis et al 2007; Eser et al 2004). Studies are supporting the concept of a new steady state at 16-24 months after injury, especially for bone metabolic process (Bauman WA 1997; Demirel et al 1998; Szollar et al 1998), but BMD decreases over the years at different areas and is inversely related to the time of the injury, which means continuous bone loss beyond the first two years after the injury (Coupaud et al 2009; Dionyssiatis et al 2008; Eser et al 2004) (Figure 2). The role played by factors such as race or gender of patients is not yet clear documented, but studies indicated more loss in women than men (Garland et al 2001; Weiss 2003). Loss of bone is closing fracture threshold from 1 to 5 years after injury (Szollar et al 1998) and risk factors for fractures after spinal cord injury are gender (women are more at risk than men), age and duration of injury (increasing age and duration of injury increases the risk of fracture with a statistically significant increase in 10 years after injury), the type of injury (complete SCI subjects have more fractures than incomplete), low body mass index (BMI) and low bone density in the tibia (Garland et al 2004; Garland et al 1992; Lazo et al 2001).

Management of bone loss in spinal cord injury

The effect of standing in bone after SCI has been investigated by many researchers. A beneficial effect on bone mass using passive mechanical loading has been shown on preservation of bone mass in the region of the femoral shaft, but not at the proximal hip of standing and non-standing patients and relatively better-preserved densities in patients standing with braces than in those using a standing frame or standing wheelchair (Goemaere et al 1994). A slower rate of bone loss in paraplegic subjects who did standing was expressed in a prospective study of 19 patients in acute SCI phase participated in early standing training program showed benefits concerning the reduction of cancellous bone loss compared to immobilized subjects (de Bruin et al 1999; Frey-Rindova et al 2000), while no correlation for passive standing-training to bone status was found in another p QCT study (Eser et al 2005). Protection afforded by standing in the femoral diaphysis stands in contrast with the loss of bone in the proximal femur. This suggests that the transmission of forces through trabecular and cortical bone varies; so the less effective strain for the initiation of bone remodelling reaches faster cortical bone (Frost 1992; Frost 2001; Frost 2003). Others also supported the concept of different strain thresholds bone remodelling control (Gutin and Kasper 1992; LeBlanc et al 2007; Smith et al 2009). There is level 2 evidence (from 1 non-randomized prospective controlled trial) that Functional Electrical Stimulation (FES) - cycling did not improve or maintain bone at the tibial midshaft in the acute phase (Eser et al 2003). Moreover, there is level 4 evidence (from 1 pre-post study) that 6 months of FES cycle ergometry increased regional lower extremity BMD over areas stimulated (Chen et al 2005). Body weight supported treadmill

training (BWSTT) did not alter the expected pattern of change in bone biochemical markers over time and bone density at fracture-prone sites (Giangregorio et al 2009).

Calcitonin in varying doses and methods of administration has given variable results in paraplegia (preferred dosage regimen, treatment duration, and administration route for adequate efficacy in SCI patients' remains unclear) (Chantraine et al 1979a; Minaire P 1987). Likewise, the outcome using bisphosphonates has been variable. Etidronate produced long-term benefit in lower limb bone mineral density (BMD) in selected walking SCI patients (Roux et al 1998); whereas tiludronate appeared effective in reducing bone resorption and preserving bone mass in a histomorphometric study in 20 paraplegic patients (Chappard et al 1995). Intravenous pamidronate has been shown to attenuate bone loss in SCI and normalize serum calcium in immobilization hypercalcemia (Bauman et al 2005). Alendronate (1000 times more potent than etidronate), in an open observational study, reversed BMD loss in men with established SCI increased both axial and trabecular bone density and has proven efficacy and safety in men treated for osteoporosis, prevents hypercalciuria and bone loss after bed rest and lower leg fracture (Moran de Brito et al 2005; Zehnder et al 2004). Six months after using zoledronic acid in the treatment group BMD showed differences in the response to treatment between the mixed trabecular/ cortical regions (narrow neck and intertrochanteric) and the purely cortical shaft. With respect to cross-sectional geometry, bone cross-sectional area and sectional modulus (indices of resistance to axial and bending loads, where higher values would indicate a positive effect of treatment) increased at the hip and buckling ratio (an index of the instability of thin-walled cross sections, where lower values would suggest that the treatment is improving stability) decreased consistent with improved bone outcomes; at 12 months, narrow-neck femur values declined and intertrochanteric and femoral shaft BMD was maintained vs. placebo group which showed a decrease in bone outcomes and an increase in buckling ratio at the hip at 6 and 12 months, while with respect to bone prevention 4 mg i.v. were effective and well-tolerated to prevent BMD loss at the total hip and trochanter for up to 12 months following SCI (Bubbear et al; Shapiro et al 2007).

Physiopathology of bone loss in Multiple Sclerosis

Reduced mobility has been implicated as an important factor in bone loss in patients suffering from multiple sclerosis (MS) and it seems to greatly influence the BMD of the femur. However, the high proportion of ambulatory patients with bone loss suggest additional non-mechanical factors (Cosman et al 1998). There is a high incidence of vitamin D deficiency in MS patients and is determined by levels of 25-hydroxy vitamin D <20ng/ml (Nieves et al 1994). The reasons might be due to a combination of low dietary vitamin D intake and avoiding of sun exposure, and that because of MS symptoms may worsen after sun exposure (fatigue-related heat) leading these patients to avoid sun. Low testosterone alone in these populations does not explain bone loss and no clear effect of smoking or alcohol abuse to decreased bone mass could be established (Weinstock-Guttman et al 2004). Glucocorticoid (GC)-induced osteoporosis (OP-GC) is the main type of secondary osteoporosis (Canalis et al 2004; Canalis et al 2007; Lakatos et al 2000; Mazziotti et al 2006; Schwid et al 1996; Shuhaibar et al 2009). The mechanism is that excess GC causes a rapid and significant damage to bone quality. Now days we know that GCs act direct on bone mainly to the stromalosteoblastic lineage and at high concentrations alter differentiation, survival, and function of them causing a shift from osteoblastic to

adipocytic differentiation of precursors; inducing apoptosis of mature osteoblasts; and inhibition of synthesis and secretion of bone components (Manolagas 2000; Pereira et al 2002). Finally, GCs promote osteoclasts and stimulate bone resorption (Weinstein et al 2002). The mechanisms of GCs action in bone has been studied extensively. In patients receiving chronic per os GC, bone loss is admitted rapidly and is evident within 6 or even 3 months (Cosman et al 1998). A study investigated the effect of intravenously (i.v.) administration of glucocorticoids in MS patients found no clear effect on bone loss: on the contrary they reported an increase in BMD of the lumbar spine (Schwid et al 1996). Prolonged treatment with glucocorticoids results in increased risk of fractures, evident at 3 months, regardless of changes in BMD. High dose, short-term i.v. treatment with GCs leads directly to reduction of bone formation and increased bone resorption, as indicated by markers of bone turnover (De Vries et al 2007; Van Staa et al 2000). In the study of Zorzon et al osteopenia not osteoporosis was significantly more frequent in patients with MS compared with controls, especially in women who received high dose methylprednisolone pulses (HDMP) in relapses period making important the regularly monitoring of BMD in these patients. The authors concluded that disability and the subsequent immobilization osteoporosis is the more serious factor in this group and treatment with repeated HDMP pulses did not cause osteoporosis in MS subjects followed-up for almost 8 years unlike chronic corticosteroid therapy which induces osteoporosis and/or recovery of BMD is permitted without permanent skeletal damage (Zorzon et al 2005). The lack of physical activity exacerbates osteoporosis.

Management of bone loss in multiple sclerosis

All MS patients should be considered high risk for osteoporosis. Prevention with calcium rich foods and dietary supplements containing vitamin D and antiosteoporotic drugs is necessary for these patients. Particular attention should be paid to transfers and falls prevention in this population to prevent fractures which occur easily and heal slowly (Cattaneo et al 2007). Subjects with multiple sclerosis are not the best candidates for bisphosphonates because of swallowing difficulties. Selective Estrogen Receptor Modulators (SERMs) and strontium ranelate could be an alternative for effective treatment of MS-related osteopenia-osteoporosis.

Conclusions

Despite the fact that the design of studies is mostly cross-sectional, it is clear that spinal cord injury and multiple sclerosis induce bone loss. The neurogenic factor seems to co-exist as an influential regulator in disabilities related bone loss during the years. Understanding the central control of bones is the cornerstone to develop treatments for neurological osteoporosis in the future.

References

- Bauman WA GD, Schwartz E. 1997. Calcium metabolism and osteoporosis in individuals with spinal cord injury. *Topics in Spinal Cord Injury Rehabilitation* 2:84-96.
- Bauman WA, Wecht JM, Kirshblum S, Spungen AM, Morrison N, Cirnigliaro C, Schwartz E. 2005. Effect of pamidronate administration on bone in patients with acute spinal cord injury. *Journal of Rehabilitation Research and Development* 42(3):305-13.

- Bubbear JS, Gall A, Middleton FR, Ferguson-Pell M, Swaminathan R, Keen RW. Early treatment with zoledronic acid prevents bone loss at the hip following acute spinal cord injury. *Osteoporosis International* 22(1):271-9.
- Canalis E, Bilezikian JP, Angeli A, Giustina A. 2004. Perspectives on glucocorticoid-induced osteoporosis. *Bone* 34(4):593-8.
- Canalis E, Mazziotti G, Giustina A, Bilezikian JP. 2007. Glucocorticoid-induced osteoporosis: pathophysiology and therapy. *Osteoporosis International* 18(10):1319-28.
- Cattaneo D, Jonsdottir J, Zocchi M, Regola A. 2007. Effects of balance exercises on people with multiple sclerosis: a pilot study. *Clinical Rehabilitation* 21(9):771-81.
- Chantraine A. 1978. Actual concept of osteoporosis in paraplegia. *Paraplegia* 16(1):51-8.
- Chantraine A, Heynen G, Franchimont P. 1979a. Bone metabolism, parathyroid hormone, and calcitonin in paraplegia. *Calcified Tissue International* 27(3):199-204.
- Chantraine A, van Ouwenaller C, Hachen HJ, Schinas P. 1979b. Intra-medullary pressure and intra-osseous phlebography in paraplegia. *Paraplegia* 17(4):391-9.
- Chappard D, Minaire P, Privat C, Berard E, Mendoza-Sarmiento J, Tournebise H, Basle MF, Audran M, Rebel A, Picot C, et al. 1995. Effects of tiludronate on bone loss in paraplegic patients. *Journal of Bone and Mineral Research* 10(1):112-8.
- Chen SC, Lai CH, Chan WP, Huang MH, Tsai HW, Chen JJ. 2005. Increases in bone mineral density after functional electrical stimulation cycling exercises in spinal cord injured patients. *Disability and Rehabilitation* 27(22):1337-41.
- Cosman F, Nieves J, Komar L, Ferrer G, Herbert J, Formica C, Shen V, Lindsay R. 1998. Fracture history and bone loss in patients with MS. *Neurology* 51(4):1161-5.
- Coupaud S, McLean AN, Allan DB. 2009. Role of peripheral quantitative computed tomography in identifying disuse osteoporosis in paraplegia. *Skeletal Radiology* 38(10):989-95.
- Dauty M, Perrouin Verbe B, Maugars Y, Dubois C, Mathe JF. 2000. Supralesional and sublesional bone mineral density in spinal cord-injured patients. *Bone* 27(2):305-9.
- de Bruin ED, Frey-Rindova P, Herzog RE, Dietz V, Dambacher MA, Stussi E. 1999. Changes of tibia bone properties after spinal cord injury: effects of early intervention. *Archives of Physical Medicine and Rehabilitation* 80(2):214-20.

- De Vries F, Bracke M, Leufkens HG, Lammers JW, Cooper C, Van Staa TP. 2007. Fracture risk with intermittent high-dose oral glucocorticoid therapy. *Arthritis & Rheumatism* 56(1):208-14.
- Demirel G, Yilmaz H, Paker N, Onel S. 1998. Osteoporosis after spinal cord injury. *Spinal Cord* 36(12):822-5.
- Dionyssiatis Y. 2009. Bone loss in paraplegia: A diagnostic and therapeutic protocol. *Osteoporosis International* 20(Suppl 1):S23-S176.
- Dionyssiatis Y, Petropoulou K, Rapidi CA, Papagelopoulos P, Papaioannou N, Galanos A, Papadaki P, Lyritis GP. 2008. Body composition in paraplegic men. *Journal of Clinical Densitometry* 11(3):437-43.
- Dionyssiatis Y, Trovas G, Galanos A, Raptou P, Papaioannou N, Papagelopoulos P, Petropoulou K, Lyritis GP. 2007. Bone loss and mechanical properties of tibia in spinal cord injured men. *Journal of Musculoskeletal and Neuronal Interactions* 7(1):62-8.
- Eser P, de Bruin ED, Telley I, Lechner HE, Knecht H, Stussi E. 2003. Effect of electrical stimulation-induced cycling on bone mineral density in spinal cord-injured patients. *European Journal of Clinical Investigation* 33(5):412-9.
- Eser P, Frotzler A, Zehnder Y, Schiessl H, Denoth J. 2005. Assessment of anthropometric, systemic, and lifestyle factors influencing bone status in the legs of spinal cord injured individuals. *Osteoporosis International* 16(1):26-34.
- Eser P, Frotzler A, Zehnder Y, Wick L, Knecht H, Denoth J, Schiessl H. 2004. Relationship between the duration of paralysis and bone structure: a pQCT study of spinal cord injured individuals. *Bone* 34(5):869-80.
- Frey-Rindova P, de Bruin ED, Stussi E, Dambacher MA, Dietz V. 2000. Bone mineral density in upper and lower extremities during 12 months after spinal cord injury measured by peripheral quantitative computed tomography. *Spinal Cord* 38(1):26-32.
- Frost HM. 1992. Perspectives: bone's mechanical usage windows. *Bone and Mineral* 19(3):257-71.
- Frost HM. 2001. From Wolff's law to the Utah paradigm: insights about bone physiology and its clinical applications. *Anatomical Record* 262(4):398-419.
- Frost HM. 2003. Bone's mechanostat: a 2003 update. *The anatomical record. Part A, Discoveries in molecular, cellular, and evolutionary biology* 275(2):1081-101.
- Garland DE, Adkins RH, Kushwaha V, Stewart C. 2004. Risk factors for osteoporosis at the knee in the spinal cord injury population. *Journal of Spinal Cord Medicine* 27(3):202-6.
- Garland DE, Adkins RH, Stewart CA, Ashford R, Vigil D. 2001. Regional osteoporosis in women who have a complete spinal cord injury. *Journal of Bone and Joint Surgery* 83-A(8):1195-200.

- Garland DE, Stewart CA, Adkins RH, Hu SS, Rosen C, Liotta FJ, Weinstein DA. 1992. Osteoporosis after spinal cord injury. *Journal of Orthopaedic Research* 10(3):371-8.
- Giangregorio LM, Thabane L, Debeer J, Farrauto L, McCartney N, Adachi JD, Papaioannou A. 2009. Body weight-supported treadmill training for patients with hip fracture: a feasibility study. *Archives of Physical Medicine and Rehabilitation* 90(12):2125-30.
- Goemaere S, Van Laere M, De Neve P, Kaufman JM. 1994. Bone mineral status in paraplegic patients who do or do not perform standing. *Osteoporosis International* 4(3):138-43.
- Gutin B, Kasper MJ. 1992. Can vigorous exercise play a role in osteoporosis prevention? A review. *Osteoporosis International* 2(2):55-69.
- Jiang SD, Jiang LS, Dai LY. 2007. Changes in bone mass, bone structure, bone biomechanical properties, and bone metabolism after spinal cord injury: a 6-month longitudinal study in growing rats. *Calcified Tissue International* 80(3):167-75.
- Kiratli BJ, Smith AE, Nauenberg T, Kallfelz CF, Perikash I. 2000. Bone mineral and geometric changes through the femur with immobilization due to spinal cord injury. *Journal of Rehabilitation Research and Development* 37(2):225-33.
- Lakatos P, Nagy Z, Kiss L, Horvath C, Takacs I, Foldes J, Speer G, Bossanyi A. 2000. Prevention of corticosteroid-induced osteoporosis by alfacalcidol. *Zeitschrift fur Rheumatologie* 59 Suppl 1:48-52.
- Lazo MG, Shirazi P, Sam M, Giobbie-Hurder A, Blacconiere MJ, Muppidi M. 2001. Osteoporosis and risk of fracture in men with spinal cord injury. *Spinal Cord* 39(4):208-14.
- LeBlanc AD, Spector ER, Evans HJ, Sibonga JD. 2007. Skeletal responses to space flight and the bed rest analog: a review. *Journal of Musculoskeletal and Neuronal Interactions* 7(1):33-47.
- Maimoun L, Fattal C, Micallef JP, Peruchon E, Rabischong P. 2006. Bone loss in spinal cord-injured patients: from physiopathology to therapy. *Spinal Cord* 44(4):203-10.
- Manolagas SC. 2000. Corticosteroids and fractures: a close encounter of the third cell kind. *Journal of Bone and Mineral Research* 15(6):1001-5.
- Mazziotti G, Angeli A, Bilezikian JP, Canalis E, Giustina A. 2006. Glucocorticoid-induced osteoporosis: an update. *Trends in Endocrinology & Metabolism* 17(4):144-9.
- Minaire P ME, Levermieux J 1987. Immobilisation bone loss: preventing effect of calcitonin in several clinical models. In: C RV Christiansen, editor. *Osteoporosis*. Copenhagen: Osteo Press. p. 603-610.

- Moran de Brito CM, Battistella LR, Saito ET, Sakamoto H. 2005. Effect of alendronate on bone mineral density in spinal cord injury patients: a pilot study. *Spinal Cord* 43(6):341-8.
- Nieves J, Cosman F, Herbert J, Shen V, Lindsay R. 1994. High prevalence of vitamin D deficiency and reduced bone mass in multiple sclerosis. *Neurology* 44(9):1687-92.
- NIH. 2001. NIH Consensus Development Panel on Osteoporosis Prevention, Diagnosis, and Therapy. *JAMA* 785-795.
- Pereira RC, Delany AM, Canalis E. 2002. Effects of cortisol and bone morphogenetic protein-2 on stromal cell differentiation: correlation with CCAAT-enhancer binding protein expression. *Bone* 30(5):685-91.
- Roux C, Oriente P, Laan R, Hughes RA, Ittner J, Goemaere S, Di Munno O, Pouilles JM, Horlait S, Cortet B. 1998. Randomized trial of effect of cyclical etidronate in the prevention of corticosteroid-induced bone loss. Ciblos Study Group. *Journal of Clinical Endocrinology & Metabolism* 83(4):1128-33.
- Schwid SR, Goodman AD, Puzas JE, McDermott MP, Mattson DH. 1996. Sporadic corticosteroid pulses and osteoporosis in multiple sclerosis. *Archives of Neurology* 53(8):753-7.
- Shapiro J, Smith B, Beck T, Ballard P, Dapthary M, BrintzenhofeSzoc K, Caminis J. 2007. Treatment with zoledronic acid ameliorates negative geometric changes in the proximal femur following acute spinal cord injury. *Calcified Tissue International* 80(5):316-22.
- Shuhaibar M, McKenna MJ, Au-Yeong M, Redmond JM. 2009. Favorable effect of immunomodulator therapy on bone mineral density in multiple sclerosis. *Irish Journal of Medical Sciences* 178(1):43-5.
- Smith SM, Zwart SR, Heer MA, Baecker N, Evans HJ, Feiveson AH, Shackelford LC, Leblanc AD. 2009. Effects of artificial gravity during bed rest on bone metabolism in humans. *Journal of Applied Physiology* 107(1):47-53.
- Szollar SM, Martin EM, Sartoris DJ, Parthemore JG, Deftos LJ. 1998. Bone mineral density and indexes of bone metabolism in spinal cord injury. *American Journal of Physical Medicine & Rehabilitation* 77(1):28-35.
- Van Staa TP, Leufkens HG, Abenhaim L, Zhang B, Cooper C. 2000. Use of oral corticosteroids and risk of fractures. *Journal of Bone and Mineral Research* 15(6):993-1000.
- Weinstein RS, Chen JR, Powers CC, Stewart SA, Landes RD, Bellido T, Jilka RL, Parfitt AM, Manolagas SC. 2002. Promotion of osteoclast survival and antagonism of bisphosphonate-induced osteoclast apoptosis by glucocorticoids. *Journal of Clinical Investigation* 109(8):1041-8.


- Weinstock-Guttman B, Gallagher E, Baier M, Green L, Feichter J, Patrick K, Miller C, Wrest K, Ramanathan M. 2004. Risk of bone loss in men with multiple sclerosis. *Multiple Sclerosis* 10(2):170-5.
- Weiss D. 2003. Osteoporosis and spinal cord injury.
- WHO. 1994. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. Report of a WHO Study Group. Geneva: World Health Organization. p. 1-129.
- Wilmet E, Ismail AA, Heilporn A, Welraeds D, Bergmann P. 1995. Longitudinal study of the bone mineral content and of soft tissue composition after spinal cord section. *Paraplegia* 33(11):674-7.
- Zehnder Y, Risi S, Michel D, Knecht H, Perrelet R, Kraenzlin M, Zach GA, Lippuner K. 2004. Prevention of bone loss in paraplegics over 2 years with alendronate. *Journal of Bone and Mineral Research* 19(7):1067-74.
- Zorzon M, Zivadinov R, Locatelli L, Giuntini D, Tonic M, Bosco A, Nasuelli D, Bratina A, Tommasi MA, Rudick RA and others. 2005. Long-term effects of intravenous high dose methylprednisolone pulses on bone mineral density in patients with multiple sclerosis. *European Journal of Neurology* 12(7):550-6.

Figures and Tables

Figure 1




p QCT in the tibia of control subject 39 years old man, slices: 4%,14%,38%



p QCT in the tibia in chronic complete AIS A paraplegic man thoracic 12 NLol 24 years old, slices: 4%,14%, 38%

Figures represent peripheral quantitative computed tomography (p QCT) tibia slices in control (a) and paraplegic subject (b), (scanner XCT 3000 Stratec Medizintechnik, Pforzheim, Germany). Areas in red represent trabecular bone, while areas in grey represent fat; pQCT allows the measurements of true volumetric densities at a minimum exposure to X-rays, assess cortical and trabecular bone density separately as well as to evaluate the geometrical properties of long bones non-invasively, adapted from: Dionyssiotis Y, Trovas G, Galanos A, Raptou P, Papaioannou N, Papagelopoulos P, Lyritis GP. Bone Loss and Mechanical Properties of Distal Tibia in High and Low Level Spinal Cord Injured Men (SA331) Abstracts ASBMR, 28th Annual Meeting, 2006, Philadelphia, Pennsylvania, USA.

Figure 2



The duration of paralysis was inversely related with bone loss in spinal cord injured subjects. Exponential correlation between volumetric trabecular bone mineral density BMDtrab and duration of paralysis in high paraplegics was found to fit best.

Table: An algorithm for the screening and management of osteoporosis in subjects with spinal cord injury (should be read top to bottom starting with the left column).

| <i>Clinical examination and management of bone loss in SCI (Dionyssiotis 2009)</i> | |
|---|--|
| Medical Evaluation | Treatments |
| <ul style="list-style-type: none"> • history of the patient (co morbidities, neurologic complications, use of drugs which impair bone metabolism, alcohol, smoking and information about the level of injury, duration of paralysis, immobilization period, onset of rehabilitation, use of assistive devices and orthoses). | <ul style="list-style-type: none"> • pharmacological treatment with bisphosphonates p.os and i.v. that have been studied in patients with spinal cord injuries and had positive effects on bone parameters. • use of calcium supplements (monitoring renal function) and vitamin D. |
| <ul style="list-style-type: none"> • anthropometric parameters (age, weight, body mass index, BMI) • clinical examination (level of injury according to American Spinal Injury Association Impairment Scale, AIS) and assessment of spasticity) | <ul style="list-style-type: none"> • education on falls prevention • counselling regarding osteoporosis and related factors and identification of fractures in regions of impaired sensation. |
| <ul style="list-style-type: none"> • imaging (bone densitometry by DXA at the hip and spine, and if possible, p QCT at the the tibia or femur) | <ul style="list-style-type: none"> • physical therapy including: a) range of motion exercises, b) loading of the skeleton to reduce bone loss, d) therapeutic standing-walking with orthoses, e) passive-active cycling |
| <ul style="list-style-type: none"> • measurement of bone turnover indices in the serum (parathyroid hormone, alkaline phosphatase, calcium, vitamin D, PINP molecule, osteocalcin) and urinary excretion of 24 hour (calcium, hydroxyproline, aminoterminal (NTx) and carboxylterminal (CTx) intermolecular cross-linking domain of bone type-1 collagen), which provide a good indicator of bone resorption. | <ul style="list-style-type: none"> • dietary interventions to improve dietary intake of calcium and nutrition indices. |